

Nonfat Dry Milk (NDM) - Medium Heat

Production Definition:

Medium Heat Nonfat Dry Milk is obtained by the removal of water from pasteurized skim milk. It contains not more than 5% moisture (by weight) and not more than 1.5% milkfat (by weight) unless otherwise indicated. Nonfat Dry Milk Medium Heat for human consumption complies with all provisions of the U.S. Federal Food, Drug, and Cosmetic Act.

Typical Compositional Range (Percentag	(e)1 Microbiological Analysis
Protein34.0 – 37.0	Standard Plate Counts2<10,000/g
Lactose49.5 – 52.0	Coliform2<10/g
Fat20.6 – 1.25	Salmonellanegative
Ash8.2 – 8.6	Listeria negative
Moisture23.0 – 4.0	Coagulase-positive
	Staphylococci negative
Other Characteristics	
Whey Protein Nitrogen (WPN)	1.51 – 5.99
Scorched Particle Content2	7.5 – 15.0 mg
Solubility Index2 Titratable Acidity2	<1.2 ml
Titratable Acidity2	<0.15%
Color2	White to light cream/natural color
Flavor2	Clean and pleasing
	te inc bein 122
Ingredient Statement	LO, IIIU.
"Nonfat Dry Milk" (% milkfat) if t	the fat content is over 1.5%

Production Applications and Functionality

Prepared mixes, ice cream, confectionery, meat products

Storage & Shipping

Product should be stored and shipped in a cool, dry environment with temperatures below 80°F and relative humidity below 65%. Stocks should be rotated and utilized within 1 to 1.5 years

Packaging

Multiwall kraft bags with polyethylene inner liner or other approved closed container. (i.e. "tote bins," etc)

¹ On an "as is" basis

² Optional tests (7 CFR §58.2608)

^{*}The information contained herein is, to the best of our knowledge, correct. No warranties, expressed or implied, are made. This information is offered solely for your investigation, verification, and consideration. It is suggested that you evaluate the product prior to use in production.*